

Conic quadratic optimization - part 4

Erling D. Andersen

MOSEK ApS,
Fruebjergvej 3, Box 16,
2100 Copenhagen,
Denmark.

Email: e.d.andersen@mosek.com

Personal WWW: <http://erling.andersen.name>

Company WWW: <http://www.mosek.com>

April 11, 2008

<http://www.mosek.com>

Introduction

[Introduction](#)

[Topics](#)

[Solving conic
quadratic
optimization
problems](#)

[The conic simplex
algorithm](#)

[Smooth convex
optimization
approach](#)

[Interior point
approaches](#)

[A primal
interior-point
approach to conic
quadratic
optimization](#)

[Summary](#)

[Exercises](#)

- Discuss different solution approaches for conic quadratic optimization problems.

Solving conic quadratic optimization problems

Introduction

Solving conic
quadratic
optimization
problems

The Ben-Tal and Nemirovski linearization

What is meant by
approximation

The decomposition
Computational
results

The conic simplex
algorithm

Smooth convex
optimization
approach

Interior point
approaches

A primal
interior-point
approach to conic
quadratic
optimization

Summary

Exercises

- Can the second order cone be approximated by a polynomial number ($O(n)$) linear inequalities?
- The answer is **yes** as proved by Ben-Tal and Nemirovski [1].
 - ◆ Using an appropriate definition of approximated.
- See also the ph.d. thesis of Francois Glineur [2].

Introduction

Solving conic
quadratic
optimization
problems

The Ben-Tal and
Nemirovski
linearization

What is meant by
approximation

The decomposition
Computational
results

The conic simplex
algorithm

Smooth convex
optimization
approach

Interior point
approaches

A primal
interior-point
approach to conic
quadratic
optimization

Summary

Exercises

Definition 1 A set $\mathcal{U} \in \mathcal{R}^n$ is said to be an ε –approximation of the second-order cone \mathcal{K}_q if and only if we have

$$\mathcal{K}_q \subseteq \mathcal{U} \subseteq \mathcal{K}_q^\varepsilon = \{x \in R^n : x_1(1 + \varepsilon) \geq \|x_{2:n}\|\}$$

Main idea:

- Prove that there is ε – approximation for the 3 dimensional quadratic cone.
- Prove that any quadratic cone can be written using a number of 3 dimensional cones.

Define the set

$$\begin{aligned}\mathcal{F}^k &:= \{(r, \alpha_0, \dots, \alpha_k, \beta_0, \dots, \beta_k) \in \mathcal{R}^{2k+3} : \\ \alpha_{i+1} &= \alpha_i \cos\left(\frac{\pi}{2^i}\right) + \beta_i \sin\left(\frac{\pi}{2^i}\right), \quad i = 0, \dots, k-1, \\ \beta_{i+1} &\geq \beta_i \cos\left(\frac{\pi}{2^i}\right) - \alpha_i \sin\left(\frac{\pi}{2^i}\right), \quad i = 0, \dots, k-1, \\ -\beta_{i+1} &\geq \beta_i \cos\left(\frac{\pi}{2^i}\right) - \alpha_i \sin\left(\frac{\pi}{2^i}\right), \quad i = 0, \dots, k-1, \\ r &= \alpha_k \cos\left(\frac{\pi}{2^k}\right) + \beta_k \sin\left(\frac{\pi}{2^k}\right)\}\end{aligned}$$

Define

$$\mathcal{G}^k := \{x \in \mathcal{R}^3 : (x_1, x_2, \alpha_1, \dots, \alpha_k, x_3, \beta_1, \dots, \beta_k) \in \mathcal{F}\}.$$

Introduction

Solving conic
quadratic
optimization
problems

The Ben-Tal and
Nemirovski
linearization

What is meant by
approximation

The decomposition
Computational
results

The conic simplex
algorithm

Smooth convex
optimization
approach

Interior point
approaches

A primal
interior-point
approach to conic
quadratic
optimization

Summary

Exercises

Lemma 1 \mathcal{G}^k is an ε -approximation for K_q^3 where

$$\varepsilon = \cos \left(\frac{\pi}{2^k} \right)^{-1} - 1$$

For a proof see [2].

Quite good result because

k	$\varepsilon \leq$
2	0.5
4	0.02
8	1.0e-4
16	2.0e-9

The decomposition

Introduction

Solving conic
quadratic
optimization
problems

The Ben-Tal and
Nemirovski
linearization
What is meant by
approximation

The decomposition

Computational
results

The conic simplex
algorithm

Smooth convex
optimization
approach

Interior point
approaches

A primal
interior-point
approach to conic
quadratic
optimization

Summary

Exercises

Now

$$\mathcal{K}_q = \left\{ x \in \mathcal{R}^n : x_1^2 \geq \sum_{j=2}^n x_j^2, x_1 \geq 0 \right\}$$

Note

$$\begin{aligned} \sum_{j=2}^{\lfloor \frac{n}{2} \rfloor} x_j^2 &\leq y_l^2, \\ \sum_{j=\lceil \frac{n}{2} \rceil}^n x_j^2 &\leq y_r^2, \\ y_l^2 + y_r^2 &\leq x_1^2, \\ 0 &\leq x_1, y_l, y_r. \end{aligned}$$

is another representation of the quadratic cone.

- The largest cone has about $\frac{1}{2}n$ variables in the new representation.
- Had to introduce 2 cones and 2 variables.
- Recursive application of this idea will produce a problem with about n 3 dimensional cones and n additional variables.

Introduction

Solving conic
quadratic
optimization
problems

The Ben-Tal and
Nemirovski
linearization

What is meant by
approximation

The decomposition

Computational
results

The conic simplex
algorithm

Smooth convex
optimization
approach

Interior point
approaches

A primal
interior-point
approach to conic
quadratic
optimization

Summary

Exercises

- If each 3 dimensional cone is ε –approximated then the approximation for the big cone is

$$\prod_{l=1}^q (1 + \varepsilon) - 1.$$

where $q \approx \log_2(n)$.

- Hence using $O(1)n$ variables and $O(1)n$ linear constraints it is possible to build a ε – approximation to the quadratic cone.
- Warning: Approximate an quadratic problem, then no bound can be given on the quality of the objective value. [2].

Introduction

Solving conic
quadratic
optimization
problems

The Ben-Tal and
Nemirovski
linearization
What is meant by
approximation

The decomposition

Computational
results

The conic simplex
algorithm

Smooth convex
optimization
approach

Interior point
approaches

A primal
interior-point
approach to conic
quadratic
optimization

Summary

Exercises

- The results of Glineur suggests
 - ◆ The linearized problems are very hard for the simplex algorithm.
 - ◆ The linearized problem can be solved using an interior-point reasonably well.
 - ◆ The primal-dual conic interior-point algorithm is much better.
- An interesting application in mixed integer conic optimization is reported in [5].

The conic simplex algorithm

A primal simplex like algorithm

Introduction

Solving conic
quadratic
optimization
problems

The conic simplex
algorithm

A primal simplex like
algorithm

Smooth convex
optimization
approach

Interior point
approaches

A primal
interior-point
approach to conic
quadratic
optimization

Summary

Exercises

Observe that any

$$x \in \mathcal{K}_q = \{x \in \mathcal{R}^n : x_1 \geq \|x_{2:n}\|\}$$

can be represented by

$$x = \sum_j \lambda_j \bar{x}^j, \lambda \geq 0$$

where

$$\bar{x}^j \in \mathcal{K}_q.$$

- x can be written as a positive sum of points in the cone.
- Extreme rays are sufficient.
- Nevertheless the sum may be infinite.
- Leads to a column generation approach.

Introduction

Solving conic
quadratic
optimization
problems

The conic simplex
algorithm

A primal simplex like
algorithm

Smooth convex
optimization
approach

Interior point
approaches

A primal
interior-point
approach to conic
quadratic
optimization

Summary

Exercises

- Developed by Goldfarb in [3].
- Not available in any commercial or academic code.
- My experience is that it works poorly in practice.

Smooth convex optimization approach

The conic quadratic constraint

$$x_1 \geq \|x_{2:n}\|$$

is equivalent to

$$x_1^2 \geq \|x_{2:n}\|^2, \quad x_1 \geq 0$$

which in turn is equivalent to

$$x_1 \geq \frac{\|x_{2:n}\|^2}{x_1}, \quad x_1 > 0.$$

almost. The function

$$f(x) = \frac{\|x_{2:n}\|^2}{x_1} - x_1$$

is a smooth, twice differentiable **convex** function on its domain.

Introduction

Solving conic
quadratic
optimization
problems

The conic simplex
algorithm

Smooth convex
optimization
approach

■
Interior point
approaches

A primal
interior-point
approach to conic
quadratic
optimization

Summary

Exercises

Hence,

$$f(x) \leq 0$$

is a convex inequality.

- The quadratic cone can be handled with a convex inequality.
- Solve the problem using any algorithm/software for smooth convex optimization e.g. conopt, loqo and Minos.
- The $x_1 > 0$ constraint can(read will) cause problems for active set methods.
- A constraint of the form $x_1 > \varepsilon$ can introduce infeasibilities.

Interior point approaches

Introduction

Solving conic
quadratic
optimization
problems

The conic simplex
algorithm

Smooth convex
optimization
approach

Interior point
approaches

Interior point
methods for linear
optimization

A barrier function

The barrier problem

The central path

A primal algorithm

A primal-dual
algorithm

A primal
interior-point
approach to conic
quadratic
optimization

Summary

Exercises

The linear optimization problem:

$$\begin{array}{ll} \min & c^T x \\ \text{st} & Ax = b, \\ & x \geq 0. \end{array} \quad (1)$$

Assumptions:

- $A \in \mathcal{R}^{m \times n}$ is of full row rank.
- $\exists x^0$ such that $Ax^0 = b$ and $x^0 > 0$.

A barrier function

Introduction

Solving conic
quadratic
optimization
problems

The conic simplex
algorithm

Smooth convex
optimization
approach

Interior point
approaches

Interior point
methods for linear
optimization

A barrier function

The barrier problem

The central path

A primal algorithm

A primal-dual
algorithm

A primal
interior-point
approach to conic
quadratic
optimization

Summary

Exercises

The function

$$B(x) = -\log(x)$$

is called a *barrier* function for the cone

$$\mathcal{K}_l = \{x \in \mathcal{R} : x \geq 0\}.$$

A barrier function is any function such that

$$\lim_{x \rightarrow 0^+} B(x) = +\infty.$$

Introduction

Solving conic quadratic optimization problems

The conic simplex algorithm

Smooth convex optimization approach

Interior point approaches

Interior point methods for linear optimization

A barrier function

The barrier problem

The central path

A primal algorithm

A primal-dual algorithm

A primal interior-point approach to conic quadratic optimization

Summary

Exercises

$$\begin{array}{ll} \min & c^T x - \mu \sum_{j=1}^n \ln(x_j) \\ \text{st} & Ax = b. \end{array} \quad (2)$$

- μ is a given **positive** parameter.
- Clearly, any feasible solution to (2) is a feasible solution to (1).
- Claim: As μ goes to 0 the optimal solution to (2) converge to the true optimal solution.

Define the Lagrange function

$$L(x, y) := c^T x - \mu \sum_{j=1}^n \ln(x_j) - y^T (Ax - b)$$

then the optimality conditions to (2) are

$$\begin{aligned} \nabla_x L(x, y) &= c - \mu X^{-1} e - A^T y = 0, \\ \nabla_y L(x, y) &= -Ax + b = 0, \\ &x > 0 \end{aligned}$$

where

Introduction

Solving conic
quadratic
optimization
problems

The conic simplex
algorithm

Smooth convex
optimization
approach

Interior point
approaches

Interior point
methods for linear
optimization

A barrier function

The barrier problem

The central path

A primal algorithm

A primal-dual
algorithm

A primal
interior-point
approach to conic
quadratic
optimization

Summary

Exercises

$$X^{-1} = \text{diag}(x_1^{-1}, \dots, x_n^{-1}) = \begin{bmatrix} x_1 & 0 & \dots & 0 \\ 0 & x_2 & & \\ \vdots & & \ddots & 0 \\ 0 & & & x_n \end{bmatrix}^{-1}$$

and

$$e = (1, \dots, 1)^T.$$

Now define

$$s := \mu X^{-1}e.$$

then the optimality conditions can be written as

$$\begin{aligned} c - A^T y - s &= 0, \\ -Ax + b &= 0, \\ s - \mu X^{-1} &= 0, \end{aligned}$$

or equivalently

$$\begin{aligned} c - A^T y - s &= 0, \\ -Ax + b &= 0, \\ Xs &= \mu e. \end{aligned} \tag{3}$$

Introduction

Solving conic
quadratic
optimization
problems

The conic simplex
algorithm

Smooth convex
optimization
approach

Interior point
approaches

Interior point
methods for linear
optimization

A barrier function

The barrier problem

The central path

A primal algorithm

A primal-dual
algorithm

A primal
interior-point
approach to conic
quadratic
optimization

Summary

Exercises

Observe

$$Xs = \mu e$$

is equivalent to

$$x_j s_j = \mu.$$

The optimality conditions (3) says:

- Dual feasibility.
- Primal feasibility.
- Perturbed complementarity.

Introduction

Solving conic
quadratic
optimization
problems

The conic simplex
algorithm

Smooth convex
optimization
approach

Interior point
approaches

Interior point
methods for linear
optimization

A barrier function

The barrier problem

The central path

A primal algorithm

A primal-dual
algorithm

A primal
interior-point
approach to conic
quadratic
optimization

Summary

Exercises

Observe that

$$\begin{aligned}c^T x - b^T y &= c^T x - (Ax)^T y \\&= (c - A^T y)^T x \\&= s^T x \\&= e^T X s \\&= \mu e^T e \\&= \mu n.\end{aligned}$$

- Conclusion: Find a solution to the barrier problem (2) for μ sufficiently small using Newton's method.
- The barrier term gets rid of the inequalities!

Introduction

Solving conic
quadratic
optimization
problems

The conic simplex
algorithm

Smooth convex
optimization
approach

Interior point
approaches

Interior point
methods for linear
optimization

A barrier function

The barrier problem

The central path

A primal algorithm

A primal-dual
algorithm

A primal
interior-point
approach to conic
quadratic
optimization

Summary

Exercises

Define $(x(\mu), y(\mu), s(\mu))$ as the solution to

$$\begin{aligned}c - A^T y - s &= 0, \\ -Ax + b &= 0, \\ Xs &= \mu e.\end{aligned}$$

- $(x(\mu), y(\mu), s(\mu))$ defines a continuous curve parameterized by μ ,
- Called the *central path*.
- Algorithmic idea.
 - ◆ Follow the central path to the optimum.
 - ◆ Leads to polynomial complexity.

A primal algorithm

Introduction

Solving conic
quadratic
optimization
problems

The conic simplex
algorithm

Smooth convex
optimization
approach

Interior point
approaches

Interior point
methods for linear
optimization

A barrier function

The barrier problem

The central path

A primal algorithm

A primal-dual
algorithm

A primal
interior-point
approach to conic
quadratic
optimization

Summary

Exercises

Define

$$F_{\mu}^p(x, y, s) = \begin{bmatrix} c - A^T y - s \\ -Ax + b \\ s - \mu X^{-1}e \end{bmatrix}.$$

Assume we are given a (x^0, y^0, s^0) such that

$$Ax^0 = b \text{ and } x^0 > 0.$$

Also let

$$\mu^0 > 0$$

be given.

Solve

$$F_{\gamma\mu^0}^p(x, y, s) = 0 \tag{4}$$

approximately where

$$0 < \gamma < 1.$$

How to solve (4) approximately? Use Newtons method i.e.

$$\nabla F_{\gamma\mu^0}^p(x^0, y^0, s^0) \begin{bmatrix} d_x \\ d_y \\ d_s \end{bmatrix} = F_{\gamma\mu^0}^p(x^0, y^0, s^0)$$

and

$$\begin{bmatrix} x^+ \\ y^+ \\ s^+ \end{bmatrix} = \begin{bmatrix} x^0 \\ y^0 \\ s^0 \end{bmatrix} + \alpha \begin{bmatrix} d_x \\ d_y \\ d_s \end{bmatrix}$$

for suitable chosen step size $\alpha \in (0, 1]$ i.e.

$$x^+ > 0.$$

The explicit Newton equation system

$$\begin{aligned} -A^T d_y - d_s &= -(c - A^T y^0 - s^0), \\ -A d_x &= -(-A x^0 + b), \\ d_s - \gamma \mu^0 (X^0)^{-2} d_x &= -(s^0 - \gamma \mu^0 (X^0)^{-1}) \end{aligned}$$

- Easy fact:

$$A x^+ = b.$$

- For suitable chosen starting point and γ :

- ◆ $\alpha = 1$ is allowed.
- ◆ Leads to convergence of order $\frac{1}{\sqrt{n}}$.
- ◆ Polynomial complexity ($O(\sqrt{n})$ iterations).

A primal-dual algorithm

Introduction

Solving conic
quadratic
optimization
problems

The conic simplex
algorithm

Smooth convex
optimization
approach

Interior point
approaches

Interior point
methods for linear
optimization

A barrier function

The barrier problem

The central path

A primal algorithm

**A primal-dual
algorithm**

A primal
interior-point
approach to conic
quadratic
optimization

Summary

Exercises

Assume we are given a (x^0, y^0, s^0) such that

$$\begin{aligned} Ax^0 &= b, & x^0 &> 0, \\ A^T y^0 + s^0 &= c, & s^0 &> 0. \end{aligned}$$

i.e. a primal-dual feasible interior solution.

Define

$$F_{\mu}^{pd}(x, y, s) = \begin{bmatrix} c - A^T y - s \\ -Ax + b \\ Xs - \mu e \end{bmatrix}.$$

The primal-dual search direction

$$\nabla F_{\gamma\mu^0}^{pd}(x^0, y^0, s^0) \begin{bmatrix} d_x \\ d_y \\ d_s \end{bmatrix} = F_{\gamma\mu^0}^{pd}(x^0, y^0, s^0)$$

Or explicitly

$$\begin{aligned} -A^T d_y - d_s &= -(c - A^T y^0 - s^0), \\ -A d_x &= -(-A x^0 + b), \\ X^0 d_s + S^0 d_x &= -(X^0 s^0 - \gamma \mu^0 e) \end{aligned}$$

- (x^+, y^+, s^+) is computed the same way as in the primal algorithm
 - ◆ Except $s^+ > 0$ is required too.
- Primal and dual feasibility are preserved.
- Is the basis for all commercially available interior-point based software for linear optimization.
- Target: Generalize this algorithm to conic quadratic optimization.

A primal interior-point approach to conic quadratic optimization

Introduction

Solving conic
quadratic
optimization
problems

The conic simplex
algorithm

Smooth convex
optimization
approach

Interior point
approaches

A primal
interior-point
approach to conic
quadratic
optimization

First algorithm

The reformulated
optimality conditions

A nonsymmetric
primal-dual
interior-point
algorithm

A small example

Summary

Exercises

Primal problem:

$$\begin{array}{ll} \min & c^T x \\ \text{st} & Ax = b, \\ & x_1 - \frac{\|x_{2:n}\|^2}{x_1} \geq 0, \\ & x_1 \geq 0. \end{array} \quad (5)$$

- We assume ONE cone for simplicity.
- The barrier function:

$$B(x) = -\frac{1}{2} \left(\ln \left(x_1 - \frac{\|x_{2:n}\|^2}{x_1} \right) + \ln(x_1) \right).$$

Introduction

Solving conic
quadratic
optimization
problems

The conic simplex
algorithm

Smooth convex
optimization
approach

Interior point
approaches

A primal
interior-point
approach to conic
quadratic
optimization

First algorithm

The reformulated
optimality conditions

A nonsymmetric
primal-dual
interior-point
algorithm

A small example

Summary

Exercises

Primal barrier problem:

$$\min_{Ax} \quad c^T x - \frac{\mu}{2} \left(\ln \left(x_1 - \frac{\|x_{2:n}\|^2}{x_1} \right) + \ln(x_1) \right) = b,$$

- μ is positive parameter.
- The optimum is in the interior for $\mu > 0$.
- Claim: For $\mu \rightarrow 0$ the optimum converges to the true optimum.

Introduction

Solving conic
quadratic
optimization
problems

The conic simplex
algorithm

Smooth convex
optimization
approach

Interior point
approaches

A primal
interior-point
approach to conic
quadratic
optimization

First algorithm

The reformulated
optimality conditions

A nonsymmetric
primal-dual
interior-point
algorithm

A small example

Summary

Exercises

Optimality conditions:

$$\begin{aligned} Ax &= b, \\ c_1 - \frac{\mu}{2(x_1 - \frac{\|x_{2:n}\|^2}{x_1})} \left(1 + \frac{\|x_{2:n}\|^2}{x_1^2}\right) - \frac{\mu}{2} x_1^{-1} - a_{:1}^T y &= 0, \\ c_{2:n} + \frac{\mu}{x_1 - \frac{\|x_{2:n}\|^2}{x_1}} \frac{x_{2:n}}{x_1} - A_{:(2:n)}^T y &= 0. \end{aligned}$$

[Introduction](#)

[Solving conic
quadratic
optimization
problems](#)

[The conic simplex
algorithm](#)

[Smooth convex
optimization
approach](#)

[Interior point
approaches](#)

[A primal
interior-point
approach to conic
quadratic
optimization](#)

[First algorithm](#)

[The reformulated
optimality conditions](#)

[A nonsymmetric
primal-dual
interior-point
algorithm](#)

[A small example](#)

[Summary](#)

[Exercises](#)

- Solve optimality conditions using Newton's method.
- While decreasing μ .
- Given the right assumptions it leads to polynomial complexity.
- Does not exploit primal-dual structure.
- A primal-dual algorithm should work better.

The reformulated optimality conditions

Introduction

Solving conic
quadratic
optimization
problems

The conic simplex
algorithm

Smooth convex
optimization
approach

Interior point
approaches

A primal
interior-point
approach to conic
quadratic
optimization

First algorithm

**The reformulated
optimality conditions**

A nonsymmetric
primal-dual
interior-point
algorithm

A small example

Summary

Exercises

Define:

$$\begin{aligned}\bar{y} &:= \frac{\mu}{2(x_1 - \frac{\|x_{2:n}\|^2}{x_1})}, \\ s_1 &:= \frac{\mu}{2x_1}.\end{aligned}$$

Reformulated system:

$$\begin{aligned}Ax &= b, \\ c_1 - \bar{y}(1 + \frac{\|x_{2:n}\|^2}{x_1^2}) - a_{:1}^T y - s_1 &= 0, \\ c_{2:n} + \bar{y} \frac{x_{2:n}}{x_1} - A_{:(2:n)}^T y &= 0, \\ \bar{y}(x_1 - \frac{\|x_{2:n}\|^2}{x_1}) &= \frac{\mu}{2}, \\ x_1 s_1 &= \frac{\mu}{2}.\end{aligned}$$

Observations:

- Is perturbed KKT system to (5).
- \bar{y} and s_1 are dual multipliers.

A nonsymmetric primal-dual interior-point algorithm

Introduction

Solving conic
quadratic
optimization
problems

The conic simplex
algorithm

Smooth convex
optimization
approach

Interior point
approaches

A primal
interior-point
approach to conic
quadratic
optimization

First algorithm

The reformulated
optimality conditions

**A nonsymmetric
primal-dual
interior-point
algorithm**

A small example

Summary

Exercises

- Solve system using Newton's method.
- While decreasing perturbation.
- Authors suggesting this idea:
 - ◆ McCormick,
 - ◆ Vial, Vial and Anstreicher,
 - ◆ Andersen and Ye,
 - ◆ Vanderbei.

A small example

Introduction

Solving conic
quadratic
optimization
problems

The conic simplex
algorithm

Smooth convex
optimization
approach

Interior point
approaches

A primal
interior-point
approach to conic
quadratic
optimization

First algorithm

The reformulated
optimality conditions

A nonsymmetric
primal-dual
interior-point
algorithm

A small example

Summary

Exercises

Study

$$\begin{array}{ll} \min & x_1 \\ \text{s.t.} & \frac{x_2^2}{x_1} \leq x_1, \\ & x_1 \geq 0. \end{array}$$

Perturbed primal-dual system:

$$\begin{aligned} 1 - \bar{y}\left(1 + \frac{x_2^2}{x_1}\right) - s_1 &= 0, \\ \bar{y} \frac{x_2}{x_1} &= 0, \\ \bar{y}\left(x_1 - \frac{x_2^2}{x_1}\right) &= \frac{\mu}{2}, \\ x_1 s_1 &= \frac{\mu}{2}. \end{aligned}$$

Observations:

- One solution

$$(x_1, x_2, \bar{y}, s_1) = (\epsilon, \beta\epsilon, 0, 1)$$

where $\beta \in (0, 1)$. Is not strictly complementarity.

- The nonsymmetric primal-dual algorithm converge to

$$(x_1, x_2, \bar{y}, s_1) = (\epsilon, \beta\epsilon, 0.5, 0.5)$$

for “bad” starting points. Point is strictly complementarity.

- See also the discussion of this example in [4].

Introduction

Solving conic
quadratic
optimization
problems

The conic simplex
algorithm

Smooth convex
optimization
approach

Interior point
approaches

A primal
interior-point
approach to conic
quadratic
optimization

First algorithm

The reformulated
optimality conditions

A nonsymmetric
primal-dual
interior-point
algorithm

A small example

Summary

Exercises

- Implies dual feasibility is not achieved.
- Computational results indicates:
 - ◆ It works for nice problems.
 - ◆ Computes primal optimal solution.
 - ◆ Has severe problems with dual feasibility.

Summary

Introduction

Solving conic
quadratic
optimization
problems

The conic simplex
algorithm

Smooth convex
optimization
approach

Interior point
approaches

A primal
interior-point
approach to conic
quadratic
optimization

Summary

References

Exercises

- Shown that conic quadratic problems can be approximated by a linear problem.
 - ◆ Is inefficient in most applications.
 - ◆ Generates hard problems for the simplex methods.
- Outline the conic simplex algorithm.
 - ◆ But has not proven to be efficient in practice.
- Discussed a couple of interior-point approaches.
 - ◆ Primal interior-point methods works well at least in theory.

Introduction

Solving conic
quadratic
optimization
problems

The conic simplex
algorithm

Smooth convex
optimization
approach

Interior point
approaches

A primal
interior-point
approach to conic
quadratic
optimization

Summary

References

Exercises

- [1] A. Ben-Tal and A. Nemirovski. On polyhedral approximations of the second-order cone. *Math. Oper. Res.*, 26(2):193–205, 2001.
- [2] F. Glineur. *Topics in Convex Optimization: Interior-Point Methods, Conic Duality and Approximations*. PhD thesis, Service de Mathématique et de Recherche Opérationnelle, Faculté Polytechnique de Mons, Rue de Houdain, 9, B-7000 Mons, Belgium, 2001.
- [3] D. Goldfarb. The simplex method for conic programming. Technical Report "TR-2002-05", IOER, Columbia University, New York, 2002.
- [4] Robert J. Vanderbei and Hande Yurttan. Using LOQO to solve second-order cone programming problems . Technical Report SOR-98-9, Department of Civil engineering and Operations Research, Princeton University, 1998.
- [5] Juan Pablo Vielma, Shabbir Ahmed, and George L. Nemhauser. A lifted linear programming branch-and-bound algorithm for mixed integer conic quadratic programs. Technical report, Georgia Institute of Technology, 2007.

Exercises

1. Prove that the function

$$f(x) = \frac{\|x_{2:n}\|^2}{x_1} - x_1$$

- is convex on its domain.
2. Consider the problem

$$\begin{array}{ll} \min & x_1 \\ \text{st} & x_1^2 + x_2^2 \leq 1. \end{array}$$

- Rewrite the problem as conic quadratic optimization problem.
 - Specify an linear relaxation that is $\varepsilon = 1.0e - 3$ accurate.
 - Solve the linear relaxation using you favorite software.
3. Implement the primal-dual interior-point algorithm for linear problems in MATLAB.